

Figure: 30 TAC §298.480(12)

United States Geological Survey Gage 08103800, Lampasas River near Kempner

Season	Subsistence	Hydrologic Condition	Base	Dry Condition Seasonal Pulse	Average Condition Seasonal Pulse	Wet Condition Seasonal Pulse
Winter	10 cfs	Dry	18 cfs	1 per season Trigger: 78 cfs Volume: 430 af Duration: 8 days	3 per season Trigger: 78 cfs Volume: 430 af Duration: 8 days	2 per season Trigger: 190 cfs Volume: 1,150 af Duration: 11 days
		Average	27 cfs			
		Wet	39 cfs			
Spring	10 cfs	Dry	21 cfs	1 per season Trigger: 780 cfs Volume: 4,020 af Duration: 13 days	3 per season Trigger: 780 cfs Volume: 4,020 af Duration: 13 days	2 per season Trigger: 1,310 cfs Volume: 6,860 af Duration: 16 days
		Average	29 cfs			
		Wet	43 cfs			
Summer	10 cfs	Dry	16 cfs	1 per season Trigger: 77 cfs Volume: 270 af Duration: 4 days	3 per season Trigger: 77 cfs Volume: 270 af Duration: 4 days	2 per season Trigger: 190 cfs Volume: 680 af Duration: 6 days
		Average	23 cfs			
		Wet	32 cfs			

cfs = cubic feet per second

af = acre-feet

N/A = not applicable