

Figure: 30 TAC §298.480(16)

United States Geological Survey Gage 08110500, Navasota River near Easterly

Season	Subsistence	Hydrologic Condition	Base	Dry Condition Seasonal Pulse	Average Condition Seasonal Pulse	Wet Condition Seasonal Pulse
Winter	1 cfs	Dry	9 cfs	1 per season Trigger: 260 cfs Volume: 1,610 af Duration: 9 days	3 per season Trigger: 260 cfs Volume: 1,610 af Duration: 9 days	2 per season Trigger: 800 cfs Volume: 5,440 af Duration: 12 days
		Average	14 cfs			
		Wet	23 cfs			
Spring	1 cfs	Dry	10 cfs	1 per season Trigger: 720 cfs Volume: 4,590 af Duration: 11 days	3 per season Trigger: 720 cfs Volume: 4,590 af Duration: 11 days	2 per season Trigger: 1,340 cfs Volume: 8,990 af Duration: 13 days
		Average	19 cfs			
		Wet	29 cfs			
Summer	1 cfs	Dry	3 cfs	N/A	N/A	2 per season Trigger: 49 cfs Volume: 220 af Duration: 5 days
		Average	8 cfs			
		Wet	16 cfs			

Cfs = cubic feet per second

af = acre-feet

N/A = not applicable